

A model for hydrophobic protrusions on peripheral membrane proteins

Edvin Fuglebakk^{1,2☐*}, Nathalie Reuter^{1,2**}

1 Computational Biology Unit, University of Bergen, Pb7803, 5020 Bergen, Norway

2 Department of Molecular Biology, University of Bergen, Pb7803, 5020 Bergen, Norway

☐Current Address: Institute of Marine Research, Norway

* Edvin.Fuglebakk@hi.no ** Nathalie.Reuter@cbu.uib.no

Abstract

With remarkable spatial and temporal specificities, peripheral membrane proteins bind to biological membranes. They do this without compromising solubility of the protein, and their binding sites are not easily distinguished. Prototypical peripheral membrane binding sites display a combination of patches of basic and hydrophobic amino acids that are also frequently present on other protein surfaces. The purpose of this contribution is to identify simple but essential components for membrane binding, through structural criteria that distinguish exposed hydrophobes at membrane binding sites from those that are frequently found on any protein surface. We formulate the concepts of *protruding hydrophobes* and *co-insertability* and have analysed more than 300 families of proteins that are classified as peripheral membrane binders. We find that this structural motif strongly discriminates the surfaces of membrane-binding and non-binding proteins. Our model constitutes a novel formulation of a structural pattern for membrane recognition and emphasizes the importance of subtle structural properties of hydrophobic membrane binding sites.

Author summary

Peripheral membrane proteins bind cellular membranes transiently, and are otherwise soluble proteins. As the interaction between proteins and membranes happen at cellular interfaces, they are naturally involved in important interfacial processes, such as recognition, signalling and trafficking. Commonly their binding sites are also soluble, and their binding mechanisms are subtle and hard to recognize. This complicates our conceptual and quantitative models of how they function, and makes binding site detection difficult. It is therefore of great interest to discover traits that are common between these binding sites, and that distinguishes them from other protein surfaces. In this work we identify simple and general structural features that facilitates membrane recognition by soluble proteins. We find that these motifs are highly over-represented on peripheral membrane proteins.

Introduction

Biological membranes are ancient and crucial components in the organisation of life. Not only do they define the boundaries of cells and organelles, but they are central to a myriad of protein-protein and protein-lipid interactions instrumental in numerous pathways [1–5]. Besides the embedded transmembrane proteins and receptors, a number of soluble proteins interact transiently with the surface of cellular and organellar membranes achieving remarkable spatial and temporal specificities. These proteins are referred to as peripheral proteins and their membrane-binding site as interfacial binding site or IBS. ~~Peripheral proteins include well-known~~ Some peripheral proteins contain lipid-binding domains such as C2 and FYVE of well-known classes, such as C2-domains and FYVE-domains that confer larger proteins the ability to bind membranes [6, 7]. Many other proteins such as lipid-processing enzymes, endogenous or secreted by pathogens are also included in this definition.

Unlike protein-protein or protein-ligand interactions, interfacial binding sites of peripheral proteins are poorly characterized in terms of amino acid composition and structural patterns. Embedded and transmembrane proteins contain well defined regions of hydrophobic surface, clearly identifying their membrane interacting segments. This is ~~seldomly~~ seldom the case for peripheral membrane proteins. Currently the prototypical peripheral membrane binding site is described as displaying a combination

of basic and hydrophobic amino acids. Attempts to characterize the energetics of membrane binding has mostly focused on electrostatic complementarity of peripheral proteins with the charged surfaces of membrane [8], rather than on the desolvation of hydrophobes which is more difficult to isolate in theoretical treatments. Nevertheless the predictive power of implicit membrane models in the prediction of membrane binding sites is a strong indication of the importance of the hydrophobic effect [9] in peripheral membrane binding. For example, Lomize ~~et al.~~ could correctly identify the experimentally known IBS of 53 peripheral peptides and proteins using a model that includes only hydrophobic, desolvation and ionization energy terms [10]. Yet in order to assert the generality of a protein-membrane binding mechanism, it is not enough to demonstrate its validity for a selected set of true positives, but it is also important to evaluate it on a control dataset.

As both small hydrophobic patches and charged residues are frequently present on protein surfaces it is challenging to distinguish membrane binding sites from the rest of the peripheral membrane proteins surface solely relying on amino acid composition. There are indications that structural considerations may allow signatures of membrane interacting hydrophobes to be defined. Terms like *hydrophobic spikes* [11,12] and *protruding loops* [10] have been used to describe membrane binding sites, prompting the idea of hydrophobes protruding from the protein globule. A close look at amphipathic helices, also motivates the concept of protruding hydrophobes. Amphipathic helices are characteristic of membrane-binding peptides and proteins. When such membrane binding helices exist, they are often found lining a protein, forming a cylindrical protrusion from the globule (e.g. ENTH domain of Epsin, PDBID: 1H0A [13], [shown in Fig 1 C and D](#)). Yet, no generalization of protruding membrane binding sites has been proposed for peripheral membrane proteins.

The purpose of this contribution is to identify structural characteristics that distinguish exposed hydrophobes at membrane binding sites from those that are frequently found on any protein surface. We propose a simple definition that formalizes the concept of protruding hydrophobes, and which can be easily computed from the protein structure. This definition allows us to systematically investigate to what extent protruding hydrophobes are found on both binding and non-membrane-binding surfaces, and to identify structural criteria for recognizing exposed hydrophobes that are likely to be important for membrane binding.

A major obstacle in developing general association models for peripheral membrane proteins is the scarcity of experimentally verified binding sites, and detailed descriptions of binding orientations. So far, computational studies on the role of hydrophobes on membrane binding sites have been based on relatively small sets of proteins with known binding sites [9,10,14]. To get around this problem and to leverage the large number of proteins for which membrane binding has been identified without a detailed characterisation of the IBS, we perform a comparative statistical analysis of ~~overall~~ protein surfaces. Given ~~a classification~~ [classifications](#) of proteins that ~~separates membrane binders from non-binders~~ [identifies membrane binders](#), we compare peripheral membrane proteins with ~~non-membrane-binding reference surfaces~~ [non-binding surfaces and reference proteins](#). With this we can extend our analysis to hundreds of protein families, rather than the few dozens for which binding sites have been partially identified by experiments.

With our simple definition of structural protrusions, we perform a statistical analysis of protruding hydrophobes in a large protein structure dataset and our results support their general role in membrane association. We find that protruding hydrophobes can be used to strongly discriminate protein surfaces involved in membrane binding from those that are not. Hydrophobes are much more frequent on protruding sites of peripheral membrane proteins than in the reference dataset, and they have a strong

tendency to cluster on positions that can simultaneously interact with the membrane. 72

Results and discussion 73




Fig 1. The definitions of *protrusions* and *co-insertable protruding hydrophobes*. Panel A shows a cartoon representation of the C2 domain of human phospholipase A₂ (PDB ID: 1RLW), and panel B shows the convex hull for the same protein. Panel C and D shows the same for an example with an aliphatic helix in the ENTH-domain (PDB ID: 1H0A). All C_α- and C_β-atoms are shown as spheres. Hydrophobes are coloured orange. The convex hull for the C_α- and C_β-atomic coordinates is shown in blue. All spheres visible on the convex-hull representation are vertex residues. *Protrusions* are defined as vertex residues with low local protein density, and shown as large spheres. *Co-insertable protruding hydrophobes* are protruding hydrophobes that are adjacent vertices of the convex hull, they are shown connected by orange lines. Small black spheres are at vertex residues that have high local density, and do therefore not meet the criteria for protrusions.

Our formalisation of the concept of protruding amino acids is illustrated in Fig 1 and described in details in *Materials and methods*. In short, it relies on firstly identifying the convex hull (in blue in Fig 1) of a coarse-grained protein model consisting of only its C_α- and C_β-atoms. We then identify amino acids located at vertices of the convex hull which intuitively are good candidates to be inserted into a membrane without inserting other residues, and without deforming the protein backbone. The model thus implicitly assumes that (1) proteins interact with the membrane without appreciable conformational change, or prior to such change and (2) that the membrane is locally flat, which is a valid approximation in most cases. In order to single out the amino acids that are most exposed to solvent, we single out amino acids (vertices) in regions of low protein density, characterized as having a low number of neighboring atoms. Solvent accessibility is a necessary condition for the hydrophobic effect to contribute to binding. In addition, regions of low local protein density are also likely to cause less disruption of lipid packing upon membrane insertion. The model was formulated based on inspection of the eight proteins included as supporting information (Table S1).

In what follows, we present results of the application of this model to characterise hydrophobic properties of protrusions in peripheral membrane proteins. We do this by

comparing a dataset of peripheral membrane proteins to a reference set of non-binding protein surface segments, and a reference set of protein surfaces not interacting typical protein structures. The reference set of non-binding surface segments (*Non-binding surfaces*) is constructed from the solvent exposed regions of trans-membrane proteins and is intended to represent structures that do not interact with the membrane, as described in-. The reference set of typical proteins (*Reference proteins*) is constructed from a protein classification excluding proteins that are classified as membrane-interacting, and is intended to represent typical protein structures, with some unknown frequency of peripheral membrane binders that are not classified as such. For consistent treatment of non-redundancy and quaternary structure modeling, these sets are compared with two variants of a set of peripheral membrane proteins (*Peripheral proteins* and *Peripheral PISA*). These data sets are described in detail in *Materials and methods*. Apart from some difference in classification, the main difference between the two set of peripheral membrane proteins is the modeling of quaternary structure, which our analysis might be sensitive to. The difference is illustrated in Fig 11.

Protruding hydrophobes in a dataset of peripheral membrane proteins




Fig 2. Hydrophobes are more common on protruding positions in peripheral proteins, than in the reference set. The plots show frequencies of hydrophobes on surface amino acids, both on protrusions (A, and C) and among all solvent exposed amino acids (B and D). Compare peripheral proteins (blue) and the non-binding surfaces (red). The horizontal axes show the mean fraction (Eq 1) of protrusions or solvent exposed amino-acids that are hydrophobic. The vertical axis shows the fraction of protein families for each set. A-D shows the comparison between the data sets *Peripheral Proteins* and *Non-binding surfaces*, and E-H the comparison between the *Peripheral PISA* and *Reference proteins*.

First we calculated the frequency of hydrophobes on protrusions in peripheral proteins-protein families and compared it to the reference dataset-datasets. In Fig 2, we observe a stark contrast between the set of peripheral proteins and the reference set (Figures non-binding surfaces (Compare figures 2 A and to 2 C). Hydrophobes occur with high frequency and in almost all families on protrusions of peripheral proteins. In the reference set on the other hand, hydrophobes on protrusions are much less tolerated, reflected by a histogram mode of zero. This While less pronounced, the distinction is also clear for the comparison with reference proteins (Compare figures 2 E to 2 G). Qualitatively, the frequency of hydrophobes on protrusions are similar in the two reference sets, but the sets of peripheral proteins differ somewhat, suggesting some sensitivity to quaternary structure modeling. For both comparisons, however, this trend is specific for protruding positions, and does not reflect a general difference in composition of surface exposed amino-acids between the data sets as shown by plots in Figures 2 B and, 2 D. Indeed, 2 F and 2 H. Indeed, if we consider the frequency of

hydrophobes on all solvent exposed residues, the distributions look quite similar with both sets having histogram modes close to 0.2. This value is in agreement with the fraction of the surface of globular proteins typically reported to be hydrophobic (for instance 0.19 in Ref. [15]). The ~~surfaces of the references set~~ *Non-binding surfaces* are in some cases very small, due to the way we ensure that these surfaces are not interacting with the membrane (see *Materials and methods*). While these small surfaces are relevant samples for calculating average frequencies, the fraction of hydrophobes on such surfaces can take more extreme values (close to zero or 1). For this reason the tails of the histograms for ~~the~~ *this* reference set are somewhat fatter than those for the peripheral membrane proteins.




Fig 3. On *Peripheral proteins*, protrusions in low density regions are more often hydrophobes, compared to the *Non-binding surfaces*. The plot shows the logarithm of the odds-ratio (Eq 10) comparing the frequency of hydrophobes on *vertex* residues in peripheral proteins and the reference set. Positive values reflect higher frequencies in the peripheral proteins. The horizontal axis shows the protein density d around the protrusion, measured as the number of C_{α} and C_{β} atoms within $1nm$. Vertex residues are all on the convex hull, but only the vertex residues with $d < 22$ are protrusions. The leftmost bar with $d < 7$ corresponds mostly to chain terminals. More precisely, the vertical axis shows $R(A, B, \hat{F}_{\text{hydrophobe}|\text{vertex} \cap l < d \leq u})$ where A denotes the *peripheral proteins*, B the *Non-binding surfaces*, l and u denote the lower and upper limits of the ranges given on the vertical axis, and d is the local protein density defined in *Materials and methods*. Error bars are 95% confidence intervals.

Given the nature of our model the differences presented in Fig 2 are naturally ascribed to two factors; the accessibility of amino acids compared to other regions of the protein (they are vertices of the convex hull) and their low local protein density d defined as the number of ~~neighbouring~~ *neighboring* C_{α} - or C_{β} -atoms (Cf. definition in *Materials and methods*). We here explore the dependence of this difference on d . In Fig 3 we show the difference between frequencies of hydrophobes in peripherals and ~~reference data sets~~ *the non-binding surfaces* for different ranges of the local protein density d . The leftmost bar ($0 \leq d \leq 6$) corresponds to chain terminals. The other bars corresponding to ranges covered by our definition of protruding residues ($7 \leq d < 22$) show that hydrophobic residues are more frequently found at vertex residues with low local protein density in the peripheral proteins. This also serves as an *a posteriori* justification for constricting our definition of protrusions to amino-acids with $d < 22$.

Assuming that the over-representation of hydrophobes on protrusions in peripheral membrane proteins stems from actual membrane binding sites, ~~one can we~~ expect those proteins to have more than one hydrophobic protrusion. We estimated the tendency of ~~each hydrophobic protrusion~~ *hydrophobic protrusions* to be *co-insertable* by calculating the weighted frequency of co-insertion (Eq 9) (Cf *Materials and methods*) for ~~both all~~ datasets (Fig 4). We note that peripheral membrane proteins do indeed tend to have




Fig 4. The protruding hydrophobes tend to be co-insertable in the peripheral proteins. The tendency for protrusions to be co-insertable is quantified by the weighted frequency of co-insertion (Eq 9), and is compared between each data set and a null model using the odds ratio (Eq 10). Positive values reflect higher frequencies of co-insertion than in the null model. More precisely, we show the comparisons $R\left(\text{set}, \text{null}, \hat{F}_{\text{one,both}}^{\text{pair}}\right)$, where *set* represents the set of peripheral proteins (blue) and the reference set (red), and *null* represent their respective null models where hydrophobes have been relocated randomly among protrusions as described in *Materials and methods*. Error bars are 95% confidence intervals. Panel A shows the comparison between the data sets *Peripheral Proteins* and *Non-binding surfaces*, and B the comparison between the *Peripheral PISA* and *Reference proteins*.

hydrophobes on co-insertable protrusions to a significantly larger extent than what would be expected from randomly scattering hydrophobes among protruding positions. This tendency is much lower for the ~~reference set~~ *Non-binding surfaces*, even when considering the extremities of the error bars, which are wide precisely because there are very few protruding hydrophobes in this set. In the Reference proteins, the analysis indicate that co-insertability is more common than in the null model, but far less so than in the Peripheral proteins.

We further explore the degree of co-insertability of the hydrophobic protrusions present in our ~~dataset of peripheral proteins and in the reference dataset~~ *datasets*. We seek to evaluate to what extent co-insertable hydrophobic protrusions can be used to discriminate likely peripheral membrane binders from other proteins. Fig 5 shows the fraction of proteins in each dataset that have at least one pair of co-insertable hydrophobic protrusions (labelled *Co-ins.*) and the fraction of proteins that have at least one *isolated* hydrophobic protrusion (i.e a protrusion that does not satisfy the criteria that define *co-insertability*). While we do see some discrimination between the data sets in the case of isolated protruding hydrophobes, the co-insertable ones prove to be very strong indicators of which proteins surfaces are membrane binding. As the coincidental occurrence of such properties increase with the size of the protein surface, we have grouped the proteins by total number of surface protrusions (regardless of hydrophobic properties). We do however see no appreciable difference between the proteins of size 0 – 25 and those of size 25 – 50. We consider the fraction in the reference ~~set-sets~~ to be a reasonable estimate of a false positive rate for predicting membrane binding function based on the presence of ~~co-insertable~~ protruding hydrophobes. ~~We find this-~~ The reference proteins (Fig 5 D-F), indicate a false positive rate to be in the range of 20% – 30%. The lack of membrane interaction is not asserted for this set, and we do expect it to contain some proteins with undetected or unclassified membrane binding. The false positive rate is around 12% for co-insertable protrusions, in both the size ranges we have analysed (see the non-binding surfaces (Fig 5) A-C), but with a smaller sample size this estimate comes with somewhat wider error bars. For the peripheral membrane proteins, ~~we estimate that around 64% and 75% of the peripheral membrane proteins in the respective groups size-groups have co-insertable protruding hydrophobes. Assuming that such motifs occur by chance at a rate no higher than it does in the reference set -, and that the over-representation is~~




Fig 5. *Co-insertable protruding hydrophobes* are common in peripheral proteins and rare in the reference set. The plot shows the occurrence of *co-insertable protruding hydrophobes* on protein surfaces. Panels A, B, D, and E show the weighted fraction (Eq 5) of proteins that have protruding hydrophobes, in the peripheral proteins (blue) and the reference sets (red). We have differentiated here between protrusions that have at least one co-insertable protruding hydrophobe (labeled “Co-ins.”), and those that have not (labeled “isolated”). The analysis is done separately for two groups of proteins according to the total number of protrusions on the protein surface ([0, 25) in panels A and D, [25, 50) in panels B and E). Panels C and F shows the frequency distribution of the total number of protruding residues (“# protrusions”) for all proteins. The selections analysed in panels A, B, D, and E are found between the dashed lines in panels C and F. Error bars in panels A, B, D, and E are 95% confidence intervals. Panels A-C shows the comparison between *Peripheral Proteins* and *Non-binding surfaces*, panels D-F, the comparison between *Peripheral PISA* and *Reference proteins*.

~~due to the membrane binding function defining the data sets, we conservatively estimate that~~ As for previous presented analysis, we see that the predictive power is weaker for the set *Peripheral PISA*. We have interpreted this as a dependence on quaternary-structure modeling, which is corroborated by Fig 11. Considering the manually curated oligomeric states two be more reliable, we expect the peripheral proteins presented in Fig 5 A-C to be a better approximation to real proteins. To get an idea of how commonly co-insertable protruding hydrophobes occur as membrane-interacting motifs, we will assume the rate of occurrence in the set *Peripheral*, and conservatively assume a frequency of occurrence on non-membrane interacting sites around 20%, which is consistent with both the extremes of the 95%-confidence intervals in the non-binding surfaces (Fig 5 A-C), and the estimate from the reference proteins (Fig 5 D-F). Even when considering that as much as 20% of co-insertable protruding hydrophobes ~~are membrane-interacting motifs for more than might not be membrane interacting, we would expect a rough estimate of around half of the peripheral-membrane-proteins we have analysed~~ analysed membrane binders to have this motif at membrane-interacting sites.

Protruding hydrophobes vs. experimentally verified membrane-binding sites

The analysis presented in Figures 3 and 5 suggests that the concepts of protruding hydrophobes and co-insertability can be used to identify membrane binding residues. Based on these results we seek to define a predictor of membrane binding sites. We define the *Likely Inserted Hydrophobe* as the protruding hydrophobe with the highest number of co-insertable protruding hydrophobes and lowest local protein density, as defined in *Materials and methods*. Fig 6 illustrates that this simple definition is able to identify binding sites on modular membrane-binding domains: C1, C2, PX, ENTH, PLA2 and FYVE. For most of these cases, the Likely Inserted Hydrophobe has in fact been experimentally ~~identified~~ indicated to contribute to membrane binding. For the




Fig 6. Protruding hydrophobes are found on the membrane binding sites of well known membrane binding domains. The figure shows the convex hull (in blue) of the C_{α} and C_{β} -atoms of selected peripheral membrane binding domains. The C_{β} -atoms of the *Likely Inserted Hydrophobe* are shown as orange spheres and C_{β} -atoms of experimentally identified membrane-binding residues as gray spheres. The Likely Inserted Hydrophobe is an amino acid that has been experimentally verified to be a membrane binding residue for A, B, D and F. For C and E the Likely Inserted Hydrophobe is located in the same area as the residues identified by experiments. **A:** C2 domain of human phospholipase A2 (PDBID: 1RLW [16]); **B:** PX domain of P40PHOX (PDBID: 1H6H [17]); **C:** snake phospholipase A2 (PDBID: 1POA [18]); **D:** C1 domain of protein kinase C delta (PDBID: 1PTR [19]); **E:** Epsin ENTH domain (PDBID: 1H0A [13]); **F:** FYVE domain of yeast vacuolar protein sorting-associated protein 27 (PDBID: 1VFY [20]).

other examples, it is clearly positioned close to the experimentally identified binding site. A more quantitative comparison between predicted and verified membrane interacting residues is complicated by the ~~absence-lack~~ of negative assertions from either methods. Experiments aiming at identifying membrane-binding sites will usually only target some of the amino acids suspected to belong to the membrane binding residues, and usually not conclude on other amino acids. To the extent non-binding amino-acids are investigated, interpretation of results in this context is also less straightforward, as non-interaction of an amino-acid does not strictly preclude it from being located close to a binding site. Similarly, the Likely Inserted Hydrophobe is by definition only one residue, and provide no negative prediction of which amino acids do not bind the membrane. We can however make a rough, but well defined, comparison by computing the angle between the vectors connecting the protein center with respectively; the mean position of the membrane interacting residues identified in experiments (\mathbf{t}_{I_e}), and the Likely Inserted Hydrophobe (\mathbf{t}_{I_p} , See Eq 11). While this comparison does not provide a quantitative evaluation of whether experimentally determined IBS and predicted residues match exactly, it allows us to separate proteins where the predicted and verified residues are “on the same side” of the protein ($\angle \mathbf{t}_{I_e} \mathbf{t}_{I_p} < 90^\circ$) from those where they are not. We show on Fig 7 such a comparison for proteins whose binding sites are experimentally determined. This is a coarse approximation to the protein orientation, which is sensitive to both protein shape, the selection of residues included in the partial binding sites, and any difference in backbone conformation between bound and unbound protein. Even so, we do expect that wrong binding site predictions should provide angles in the entire range from 0° to 180° with roughly uniform probability. But, we observe that almost all angles are sharper than 90° , indicating a reasonable agreement




Fig 7. Protruding hydrophobes predict experimentally verified binding sites. The figure shows comparisons of predicted binding residues (*the Likely Inserted Hydrophobe*) with experimentally verified binding sites for a manually curated dataset of 24 proteins (listed in Table S2). The vertical axis corresponds to values of the angle (Eq 11) comparing the two vectors connecting the center of the protein with either the predicted or known binding sites. Smaller angles imply better agreement between prediction and experiment. Asterisks (*) mark proteins where the Likely Inserted Hydrophobe is an amino acid experimentally identified to be interacting with the membrane. The grey boxplots show the distribution of angles when the known binding site residues are compared to all protruding amino acids on the protein. 1iaz is analysed in its soluble monomeric state, while it forms a transmembrane pore upon oligomerisation. The structure of the Bovine α -lactalbumin (PDBID: 1F6S) has no identified protruding hydrophobes and is marked with a cross at 180°.

with experimental data. We also observe a similar range of angles for cases where the membrane interaction of the Likely Inserted Hydrophobe has been experimentally verified (marked with asterisks (*) in Fig 7) and the cases where it has not. We would like to emphasise at this point that the Likely Inserted Hydrophobes that are not yet found to be membrane interacting might very well never have been tested. We also calculated all angles between the set of experimentally identified residues and protruding amino acids of all kinds. These results are displayed as box-plots in Fig 7. While they vary a bit between families, we note that all medians are close to 90°, confirming that the statistical expectation for protrusions in general is to have roughly equally many observations larger than and smaller than 90°. [Interestingly, the Bovine \$\alpha\$ -lactalbumin, for which we find no protruding hydrophobes, is analysed in its crystallised form while it is known to bind membranes in a molten globule state \[21\].](#)

We provide as Supporting Information the complete list of amino acids experimentally identified as being part of membrane binding sites (S1 Text, Table S2). It overlaps with the list provided by Lomize *et al.* [10], but sometimes differ in exactly which amino acids are included, as we include [indicated](#) membrane interacting residues even when they are not inserted in the hydrophobic core of the membrane.

Protruding hydrophobes on predicted membrane binding sites

The continuum-model presented by Lomize *et al.* [22] forms the basis for a systematic effort to predict binding orientations for peripheral membrane proteins. The OPM database [23] provides prediction of spatial arrangements of membrane proteins with




Fig 8. Comparing predictions based on protruding hydrophobes with the predicted IBS in the Orientation of Proteins in Membranes (OPM) database. The plots show the distributions of the median *insertion coordinate* from OPM for the *Likely Inserted Hydrophobe* in each family (measured at the C_{α} -atom). Values greater than or equal to zero correspond to atoms positioned in the hydrophobic core or at the boundary. Hence insertion coordinate values close to zero indicate agreement with OPM. Panel A (C) show data for the Likely Inserted Hydrophobes and panel B (D) for a null model of randomly selected *protruding* residues. Panel C and D show cumulative histograms (accumulated with decreasing insertion coordinates).

respect to the lipid bilayer for a selection of peripheral membrane proteins. We here investigate to what extent protruding hydrophobes are captured by the model proposed by Lomize *et al.* We identify The Likely Inserted Hydrophobe for each of the proteins in our dataset, and ~~extracts~~ extract the OPM predicted insertion coordinate of its C_{α} -atom. The *insertion coordinate* of an atom measures its depth of insertion into the hydrocarbon region of the membrane model, and is thus positive for atoms located in the hydrocarbon core and negative for atoms located on either side of the membrane including the interfacial region (Cf. *Materials and methods*). Fig 8 shows histograms of the median insertion coordinate of the Likely Inserted Hydrophobes identified in each family. A clear majority of those residues are located close to the interface of the membrane model in the OPM-predictions (Fig 8 A) and 75% of the families in the set of peripheral membrane proteins have the median insertion coordinate for the Likely Inserted Hydrophobe within a margin of 0.5 nm from the membrane. This fraction is similar to the estimated fraction of proteins that have co-insertable protruding hydrophobes (Figures 5 A and B). We allow this margin of 0.5 nm to compensate for the assumptions of rigid protein, flat membrane, and the distance between C_{α} -atoms and side-chain atoms. Fractions for other margins can be read from the cumulative histogram shown in Fig 8 C. By representing position with the insertion coordinate, we effectively project residue coordinates onto the membrane normal. We therefore do not expect surface amino acids to be uniformly distributed along the insertion coordinate axis and present control statistics for randomly chosen protruding amino acids of all hydrophobic properties (Fig 8 B and 8 D). It appears clearly that the high number of Likely Inserted Hydrophobes close to the membrane model is not an effect of it simply being more protein there.

Structure and amino acid composition at hydrophobic protrusions

280
281




Fig 9. In peripheral proteins, hydrophobic protrusions are more frequent on turns, bends and α -helices, compared to the reference set. Panel A shows the weighted number (Eq 2) of *protruding hydrophobes* associated with the different types of secondary structure elements. We have differentiated between protrusions that have at least one co-insertable protruding hydrophobe (right, labeled “Co-ins.”), and those that have not (left, labeled “Isolated”). Panel B compares the weighted frequencies (Eq 4) of hydrophobes on protruding secondary structures between the peripheral membrane proteins and the reference set, using the odds ratio (Eq 10). Positive values reflect higher frequencies in the peripheral proteins. More precisely, panel A show the values $N_{\text{hydrophobe}|\text{protrusion} \cap \text{sse}}$, and panel B the comparisons $R(A, B, \hat{F}_{\text{hydrophobe}|\text{protrusion} \cap \text{sse}})$ where A denote the peripheral proteins, B the reference set, and sse specifies the secondary structures given in the color legend. Error bars in panel B are 95% confidence intervals.

The analysis presented in Fig 3 indicates that the ability to discriminate the data sets based on the frequency of hydrophobes on protrusions gets lower as the local protein density gets higher. Local protein density of a protrusion is dependent on secondary structure elements with loops, turns and bends being those that intuitively favor low local protein density. These secondary structures typically mark a clear change in direction of the backbone trace, where the neighbouring residues ‘make way’ for the protruding hydrophobe. Fig 9 A shows which secondary structure elements the protruding hydrophobes are associated ~~to~~ with in the set of peripheral proteins. We note that loops, turns and bends are indeed abundant, but also helices and not beta-strands. Fig 9 B shows a comparison with the reference data set. We see that protruding hydrophobes on turns and bends are not only common in the peripheral membrane proteins as we saw in Fig 9 A, but they are also significantly more frequent than in the reference set. Interestingly, this is not the case for loops. A reason for this might be that turns and bends provide a rigid scaffold for exposing the hydrophobes, which would otherwise rearrange to desolvate when exposed to solvent, and thereby likely reduce the free-energy gain of membrane insertion. As the definition of *loop* here is simply absence of any of the other secondary structure definitions, we would expect this category to contain less regular, more flexible structures. We also expect this property of rigid scaffolding from amphipathic helices, which is an established motif for

282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300

membrane association. Fig 9 illustrates however that protrusions are not dominantly helices, confirming that the concept of protruding hydrophobes provides a useful generalisation for the shapes of membrane-binding sites.




Fig 10. Large aliphatic and aromatic side chains are particularly over-represented on protrusion on peripheral proteins. Panel A shows the weighted fractions (Eq 4) of hydrophobic amino acids on protrusions from peripheral proteins (blue) and from proteins in the reference set (red). In panel B, the contrast between the two sets is quantified by the odds ratio (Eq 10), so that positive values reflect higher frequencies in the set of peripheral proteins than in the reference set. More precisely the vertical axis denote $\ln R$ (peripheral, reference, $\hat{F}_{aa, protrusion}$), with aa representing each of the standard amino acids. Error bars are 95% confidence intervals.

For purposes of isolating the structural component of hydrophobic membrane association, we have until now used a dichotomous definition of hydrophobicity based on the Wimley-White scale for interfacial insertion [24] signs of free energy of transfer determined by Wimley and White [24] (leucine, isoleucine, phenylalanine, tyrosine, tryptophan, cysteine and methionine have been considered to be hydrophobic). Yet, we do expect different amino acids to have varying contributions to the free energy of binding. We have therefore also assessed the relative importance of different amino acids for discriminating between our two data sets. Fig 10 B shows the comparison of the frequencies of different hydrophobic amino acids on protrusions in the two data sets Peripheral proteins and the Non-binding surfaces. Analysis of the other two sets can be found as supplementary material. As expected, we find non-polar residues with large aliphatic or aromatic side chains to be much more frequent at the protrusions of peripheral proteins Peripheral proteins than in the reference data set set of non-binding surfaces. While the error bars in Fig 10 B are not corrected for multiple testing, the signal for the hydrophobes as a group is quite clear. They all occur as over-represented in the set of peripheral proteins Peripheral proteins and the odds-ratio is much larger for phenylalanine, leucine and tryptophan than for any of the amino-acids that are over-represented in the reference set. Non-binding surfaces. Analysis of the other two sets can be found as supplementary material. Recall that $\ln R$ (Eq 10) is symmetric around 0, so the magnitude of the bar representing phenylalanine on one end, can be directly compared to that of the bar representing threonine in the negative direction. Tyrosine on the other hand discriminates the sets poorly compared to its high hydrophobicity score in the Wimley-White scale. We consider this a possible consequence of the orientational restrictions on the binding sites

of peripheral membrane proteins. The typical orientations consistent with shallow binding, has the residue anchored above the membrane. This probably allows less freedom for the hydroxyl group of tyrosine to orient towards regions of higher water density, than it has in the peptides used for the Wimley-White experiments or in transmembrane proteins. We also note with interest that proline is among the residues that are somewhat over-represented in the set of peripheral proteins. In general, prolines are conformationally important protein components, that restricts the backbone with respect to its immediate neighbours along the peptide chain, and are therefore likely to promote local rigidity. They also serve to induce sharp changes in the backbone trace, which would facilitate solvent exposure of neighbouring side-chains, as discussed above. Specifically, they are in general frequently found on turns [25].

Comments on the protein model

The convex-hull representation presents a useful abstraction of proteins for considering surface properties for shallow interaction with an approximately flat membrane with an approximately rigid protein conformer. This enables statistical analysis of protein structures, which is prohibited by high-resolution models where model parameters and quality controls often have to be made subjectively on individual proteins. We have employed this abstraction specifically to quantify and understand aspects of the role of hydrophobes in peripheral membrane binding. In order to isolate components contributing to membrane binding, we have purposefully avoided complicating the interpretation with known important factors, such as electrostatics, conformational flexibility, and even relative hydrophobicity. For the purposes of understanding the competition and complementarity between different generic models in explaining membrane-binding, it will be necessary to take these other binding factors into account in ways that allows decomposition of their contribution. In the framework of non-energetic structural analysis presented here, it is natural to do that in terms of comparing presence, absence and location of predicted binding sites between proteins. Particularly, models of electrostatic binding are well developed and readily applicable to surface representations of rigid protein conformers. For comparison of partial binding-site indications we have developed a measure of agreement that can be applied to models of low resolution, and to compare models of different resolution. We expect this to be useful in analysing the complementarity of hydrophobic and electrostatic binding modes. While complex energetic models, or machine learning approaches, can be expected to yield high performance in predicting membrane-binding properties of a protein, the transparent kind of model presented here provides a clear interpretation of why a specific protein is predicted to bind. Apart from the obvious academic interest, this connection to expert knowledge is invaluable for interpreting automated classifications where the models can not be reliably parameterised against negative data, that is: definitely non-binding proteins. Co-localized binding-site indicators based on different generic binding models, such as hydrophobic and electrostatic models, can provide a much improved performance in such prediction while maintaining interpretability. Such decomposition of binding modes are also useful for inference or interpretation about the specificity towards interacting membranes, as electrostatic binding modes are expected in interactions with highly charged membranes, while predominantly hydrophobic interactions are more likely to be specific to membrane or lipid morphologies.

Conclusion

Protein-membrane interactions are typically studied *in vitro* or *in silico* and inference to their biological context have to carry over from greatly simplified membrane models. To make sense of such experiments and simulations, it is essential to formulate general models that explain protein association in terms of factors that are present in both model systems and the relevant *in vivo* counterpart. In pursuit of such general models for membrane recognition, we have formulated the concepts of protruding hydrophobes and co-insertability. We have analysed more than 300 families of proteins that are classified as peripheral membrane binders and identified this model to be a good fit to ~~more than at least~~ half of them, after ~~cautiously~~ correcting for the ~~small false positive rate~~ ~~false positive rates~~ estimated from the reference ~~set-sets~~ (Fig 5). The generality of the model is corroborated by three important points. Hydrophobes are clearly over-represented on the protrusions of peripheral membrane proteins (compare Fig 2 A and 2 C, and see Fig 3), they tend to locate on co-insertable protrusions (see Fig 4 and Fig 5), and protruding hydrophobes are generally positioned consistent with experimentally identified binding sites (Fig 6 and Fig 7). Amphipathic helices are already well known membrane binding motifs which our definition of protrusion is well suited to capture, whenever these are stably folded and exposed. We do however find that the majority of identified protruding hydrophobes are not helices (Fig 9 A) and that hydrophobes are also highly over-represented on protruding turns and bends (Fig 9 B). We therefore propose the concept of protruding hydrophobes as a useful generalisation upon binding motifs that are identified in terms of secondary structure.

~~Both the choice of reference set, and the choice of quaternary structure modelling comes with some assumptions. We have elaborated on these in “materials and methods”. We have also performed some checks on how sensitive our analyses are to violations of these assumptions, and found that our conclusions are robust. We present details of these analyses as Supporting Information (-).~~

Investigation of the interfacial binding sites of numerous peripheral membrane proteins has revealed the presence of hydrophobic amino acids, and of basic amino acids such as arginines and lysines. This reflect the two almost universal traits of biological membranes; their hydrophobic core and anionic surface. Yet the focus on the electrostatic component of the free energy of transfer from water to membrane - often referred to as being long-range - has overshadowed the importance of hydrophobic contribution which is sometimes referred to as being short-range. The focus on electrostatic interaction is at least in part to be attributed to the difficulties in evaluating the hydrophobic contribution as opposed to for example, the computational tractability of continuum electrostatic models. In principle the contribution of hydrophobes to membrane binding can only be determined with a rigorous treatment of the hydrophobic effect, which requires very accurate treatment of large systems involving both protein, membrane and solvent. The mere presence of hydrophobes on the protein surface is to a large extent tolerated by non-membrane-binding proteins as well, and for both hydrophobes and basic amino acids, it is challenging to determine when their presence on protein surfaces are coincidental, and when they are important for membrane binding. Moreover, amino acids on membrane binding sites are not typically strongly conserved [26], so modeling their generic binding modes is important both for relating binding sites between homologs, and for understanding how additional factors determine differences in membrane specificities. Fortunately, as evident from the results presented in this contribution, the role of hydrophobes can often be understood in much simpler terms than what is required for an exact estimate of the energetics of the hydrophobic effect, and their importance for membrane-binding can be inferred from comparative statistical analyses. The subtle considerations of protein structure encoded in our definition of protrusions, strongly distinguishes the small hydrophobic

patches on peripheral membrane proteins from those on other protein surfaces. This provides good reason to assume their importance for binding.

Materials and methods

Data sets

We ~~obtained data sets from the collection of proteins in~~ have compiled four data sets, two versions of a set of peripheral proteins, and two different reference sets:

- *Peripheral Proteins*: A set of peripheral membrane binders obtained from the OPM-database [23], using the OPM quaternary structure model.
- *Peripheral PISA*: A subset of *Peripheral Proteins* with quaternary structure predicted by PISA [27], where no domain in any protein share SCOPE-family classification [28] with domains of any other protein in the set. ~~Our set of peripheral proteins are all-~~
- *Non-binding Surfaces*: A set of protein surfaces obtained from the solvent exposed regions of transmembrane proteins.
- *Reference Proteins*: A non-redundant set of proteins from 5 SCOPE-classes, where no domain exists in OPM, and no domain in any protein share SCOPE-family classification with domains of any other protein in the set.

In our analysis *Peripheral Proteins* is always compared to *Non-binding surfaces*, and *Peripheral PISA* to *Reference Proteins*.

Peripheral Proteins are all the proteins in OPM classified as *type: Monotopic/peripheral*. While the OPM has strict criteria for inclusion, membrane binding is not asserted by experiment in all cases, and the set might contain false positives. {refer suppmat}

The ~~reference set set~~ *Non-binding surfaces* consist of fragments of transmembrane complexes. We obtained these protein fragments from all proteins classified as *type: Transmembrane* in OPM. The fragments analysed are composed of all amino acids whose C_{α} -coordinates are at least 1.5 nm from the hydrocarbon region of the membrane model (The parameter Z_{HDC} in the OPM model [29]). We rely here on membrane models positioned by the OPM, which we deem very reliable for transmembrane proteins. While the entire protein complex was considered when calculating structural properties, only the fragments meeting this distance criteria were considered in the statistical analyses. When these proteins interact with secondary membranes or interact with membranes of extremely high curvature, it is not captured by the OPM model, and the assumption that these surfaces are not interacting with membrane may be violated. We have assumed that such issues are exceptional. {refer suppmat}

We do consider the assumptions mentioned above to be conservative. Inclusion of non-membrane-binding proteins in our set of peripheral membrane proteins would likely weaken any general signal from membrane binding proteins, and inclusion of secondary membrane interactions sites in the reference set would probably inflate the number of hydrophobes on protrusions in that set.

All protein structures in these two sets are obtained by X-ray crystallography and NMR spectroscopy and we have assumed that at least the backbone coordinates are representative of the solvated state of the proteins. As the source of structural information for this database is the Protein Data Bank (PDB) [30] the relevant oligomeric state is not always determined, The curators of the OPM-database have decided on oligomer models, upon which we have relied for the sets *Peripheral Proteins*

and *Non-binding surfaces*. These are taken from PDB [31], generated by PISA [27] or obtained from literature as described by Lomize *et al.* [23]. ~~As weak protein-protein interaction interfaces may also contain exposed hydrophobic patches, we~~

Even if the solvent exposed regions of the proteins in the set *Non-binding surfaces* are extracted after relevant properties for potential membrane interaction was calculated, we are somewhat concerned that the surface constructed does not reflect all relevant properties of non-binding proteins. In addition we expect our analysis to be sensitive to ~~how protein quaternary structure is modelled~~ quaternary structure modeling, as weak protein-protein interaction interfaces may also contain exposed hydrophobic patches. As a quality control, we therefore also performed our analysis relying solely on computationally predicted quaternary structures, ~~which we provide in the Supporting Information ()~~ and complete protein structures. This is implemented in the comparison of *Peripheral PISA* and *Reference Proteins*. ~~This control reproduced qualitatively all observations that we have interpreted. In the Supporting Information we also report analysis on the sensitivity of the results to how the reference set is obtained, using a reference~~

The set *Reference Proteins* is constructed as a subset of all PDB IDs determined by X-ray crystallography, with some domain classified in SCOPe [28] in the classes *All alpha proteins* (sunid: 46456), *All beta proteins* (sunid: 48724), *Alpha and beta proteins (a+b)* (sunid: 51349), *Alpha and beta proteins (a/b)* (sunid: 53931) or *Multi domain proteins* (sunid: 56572). The exclusion of structures not determined by X-ray crystallography was done to get consistent quaternary structure predictions. All PDB IDs that has one or more domains classified in the same SCOPe-family as any domain in the OPM-database [23] was excluded from the set. This excludes not only the peripheral membrane binders, but also any transmembrane protein found in the reference set used for our primary analysis. In order to avoid redundancy, we iteratively removed proteins with domains that share SCOPe-family classification with any other domain in the set ~~based on the SCOPe-classification [28]~~, until there were no such shared classifications left. This process ensures that there is at most one representative for each SCOPe family in the set. We generated quaternary structure models using PISA [27] for all members of this set. {ref suppmat}. While this data set consist of more complete protein surfaces, it is only intended to be a reference for typical protein surfaces, we do expect it to be a mix of both membrane interacting and non-interacting proteins.

The set *Peripheral PISA* was derived from *Peripheral Proteins*, for comparability with *Reference Proteins*. All structures not determined by X-ray crystallography was excluded, and proteins with domains that share SCOPe-family classification with any other domain in the set was iteratively removed until no such shared classifications were left. Quaternary structure models were predicted using PISA. {ref suppmat}.

A few structures meeting the above criteria, were not included in the analysis for technical reasons, such as issues with formatting of PDB files. After exclusion of these cases, the final set of ~~peripheral proteins~~ *Peripheral proteins* contains 1012 protein structures classified into 326 families. The final ~~reference set~~ *set of Non-binding surfaces* contains 495 protein structures classified into 158 families. {add stats on alternative sets.}

The two set of peripheral proteins are both derived from OPM, but *Peripheral PISA* is organized in a different classification from *Peripheral Proteins*, and retains fewer structures. In addition their quaternary structures, which are not completely determined by X-ray crystallography, are modeled differently. In Figure 11, we illustrate this difference in quaternary structure by showing the difference in the number of polypeptide chains present in these protein-models in the two sets.

Based on experiments reported in available




Fig 11. Differences in number of polypeptide chains between the protein models present in the *Peripheral Proteins* (quaternary structure model from OPM) and the models in *Peripheral PISA* (quaternary structure model predicted by PISA). The difference is calculated for each of the PDB IDs occurring in both datasets. When more chains are present in the PISA models, The difference (horizontal axis) is negative.

literature [11, 21, 32, 33, 33–36, 36, 37, 37–65], we made a data set of partially identified membrane binding sites on proteins with resolved structures. This set contains membrane interacting residues of 34 protein structures, classified into 22 families. A detailed description is provided in the Supporting Information (S1 Text, Table S2).

Definitions

Structural characteristics of protein surfaces

We characterise the surface of proteins with different criteria designed to capture solvent-exposed residues, protruding residues and co-insertable protruding residues. The two latter are illustrated in Fig 1.

Exposed amino acids are defined as all amino acids that have a solvent accessible side-chain area greater than 0.2 nm², as calculated with a probe with a radius of 0.14 nm, following the procedure described in Eisenhaber *et al.* [66] using van der Waals radii reported by Bondi [67].

We identify a *protrusion* or a *protruding* residue via the calculation of the convex hull of the C_α- and C_β-coordinates of the protein. The convex hull of a set of points *S* is the smallest possible convex set containing *S*. We define *vertex* residues as residues whose C_β-atom is a vertex of this convex hull. A *protrusion* or a *protruding* residue, is defined as a *vertex* residue that also has low local protein density. For the purposes of this work, we will define the local protein density *d* of a residue, as the number of C_α- or C_β-atoms within a distance *c* of its C_β-atom. We will designate a local protein density as low, if *d* < *n*, with *n* = 22 and *c* = 1 nm. These parameters were manually chosen based on a set of six different families of peripheral membrane proteins (C2-domain, PX-domain, Discodin domain, ENTH domain, Lipoxygenases and a Bacterial Phospholipase C). A list of these proteins are provided as Supporting Information (S1 Text, Table S1).

We define two protrusions to be *co-insertable* or a *co-insertable pair*, if the straight line connecting them is an edge of the convex hull polygon.

Hydrophobic residues 550

An amino acid is defined to be *hydrophobic*, or a *hydrophobe*, if it contributes favourably to membrane interface partitioning of peptides, as determined in the Wimley-White scale for interfacial insertion [24]. These amino acids are: leucine, isoleucine, phenylalanine, tyrosine, tryptophan, cysteine and methionine. 551
552
553
554

Secondary structure 555

We use DSSP definitions [68] for protein secondary structure. DSSP codes H, G or I are reported as *helix*, DSSP codes B or E as β , DSSP code T as *bend* and DSSP code S as *turn*. All other residues are considered to be in *loops*. 556
557
558

Likely Inserted Hydrophobe 559

The *Likely Inserted Hydrophobe* is defined as the protruding hydrophobe with the largest number of co-insertable protruding hydrophobes in a protein. Ties are resolved by choosing the likely inserted hydrophobe with the smallest local protein density d . Further ties are resolved by random selection, so that each protein has exactly one Likely Inserted Hydrophobe, unless it has no protruding hydrophobes at all. 560
561
562
563
564

Insertion coordinate 565

For comparisons with OPM predictions, we define the *insertion coordinate* of atoms. This coordinate measures how deeply into the OPM membrane model an atom is inserted, and is therefore negative on the solvated side of the membrane. The membrane perimeter, where the insertion coordinate is 0, is the end of the hydrocarbon region. We identify this boundary as it is done in the model used to predict the OPM orientations, namely the planes where the volume fraction of total hydrocarbon is equal to 0.5. See Eq 2 in [29]. 566
567
568
569
570
571
572

Measures 573

Averages of residues 574

We compare protein surfaces with respect to structural and hydrophobic properties, reflected in different selection criteria and averaged over families or the entire data sets. 575
576

The mean fraction of residues having property s with respect to a reference property r in a family is: 577
578

$$\hat{f}_{s|r} = \frac{1}{|C|} \sum_{G \in C} \frac{|G_s \cap G_r|}{|G_r|} \quad (1)$$

where C is the set of proteins in a family, G is a protein, and, G_s is the set of residues on a protein meeting criteria s . Vertical bars denote size of sets. We will specify s and r according to the definitions above, using intersect notation to combine criteria when necessary. $\hat{f}_{\text{hydrophobe|protrusion} \cap \text{helix}}$, for instance, should be interpreted as the mean fraction of hydrophobes out of all protruding amino acids that are in helices. 579
580
581
582
583

We estimate weighted data set counts of amino acids with property s as: 584

$$\hat{N}_s = \sum_{C \in D} \left(\frac{1}{|C|} \sum_{G \in C} |G_s| \right) \quad (2)$$

where D is a data set, such as the set of peripheral proteins or the reference set. Similarly we quantify the weighted count of proteins that have at least one amino acid 585
586

with property s as:

$$\hat{M}_s = \sum_{C \in D} \left(\frac{1}{|C|} \sum_{G \in C} H(|G_s|) \right) \quad (3)$$

where H is the Heaviside step function. Given a property s and reference property r , we estimate the weighted fraction in a data set, $\hat{F}_{s|r}$:

$$\hat{F}_{s|r} = \frac{\hat{N}_{s \cap r}}{\hat{N}_r} \quad (4)$$

or the weighted fraction of proteins that have at least one residue with the given property s :

$$\hat{E}_s = \frac{\hat{M}_s}{|D|} \quad (5)$$

With $|D|$ being the number of families in the data set. When such fractions (Eqs. 4 or 5) are reported, we estimate 95%-confidence intervals using a normal approximation to the binomial distribution, with $|D|$ the total number of trials (Eq 5), or \hat{N}_r serving as a real-number analog to the total number of trials (Eq 4).

Averages of co-insertable pairs

To analyse co-insertable residues, we estimate weighted data set counts of co-insertable pairs of residues with property s , as:

$$\hat{N}_s^{\text{pair}} = \sum_{C \in D} \left(\frac{1}{|C|} \sum_{G \in C} |G_s^{\text{pair}}| \right) \quad (6)$$

where $|G_s^{\text{pair}}|$ are the number of co-insertable amino acids pairs with property s . For quantification of the weighted count of proteins that have at least one co-insertable pair with property s , we calculate:

$$\hat{M}_s^{\text{pair}} = \sum_{C \in D} \left(\frac{1}{|C|} \sum_{G \in C} H(|G_s^{\text{pair}}|) \right) \quad (7)$$

Considering the set of co-insertable amino acid pairs in a protein, G^{pair} , we will denote the set of pairs where at least one of the amino acids is a protruding hydrophobe as $G_{\text{one}}^{\text{pair}}$, and the set where both are protruding hydrophobes as $G_{\text{both}}^{\text{pair}}$. We will report the weighted fraction of proteins that have co-insertable protruding hydrophobes as:

$$\hat{E}_{\text{both}}^{\text{pair}} = \frac{\hat{M}_{\text{both}}^{\text{pair}}}{|D|} \quad (8)$$

and the weighted frequency of co-insertion of protruding hydrophobes as:

$$\hat{F}_{\text{both}|\text{one}}^{\text{pair}} = \frac{\hat{N}_{\text{both}}^{\text{pair}}}{\hat{N}_{\text{one}}^{\text{pair}}} \quad (9)$$

Note that $\hat{F}_{\text{both}|\text{one}}^{\text{pair}}$ estimates the conditional probability that both amino acids of a co-insertable pair are protruding hydrophobes, given that one of them is. The tendency for protruding hydrophobes to be located at co-insertable positions can then be quantified by comparing with a null model for each set. We obtain these null models by randomly reassigning the hydrophobic amino acids to other protruding locations in the same protein.

Comparison between data sets

The frequency of properties in different data sets, are compared via weighted fractions. For two data sets, A and B , we compare a certain weighted fraction \hat{F} using the odds ratio, $R(A, B, \hat{F})$:

$$R(A, B, \hat{F}) = \frac{\hat{F}^A (1 - \hat{F}^B)}{\hat{F}^B (1 - \hat{F}^A)} \tag{10}$$

where \hat{F}^A denotes the fraction $\hat{F}_{s|r}$ obtained for data set A . We will report $\ln R$, which is symmetric around 0, so that $\ln R(A, B, \hat{F}) = -\ln R(B, A, \hat{F})$. Wald 95%-confidence intervals for $\ln R$ are calculated with $\hat{N}_{s \cap r}$ and $(\hat{N}_r - \hat{N}_{s \cap r})$ serving as real number analogs for the count of successes and failures in the data sets compared. When $\hat{F}_{\text{both}|\text{one}}^{\text{pair}}$ is compared, the corresponding counts of successes and failures are $\hat{N}_{\text{both}}^{\text{pair}}$ and $\hat{N}_{\text{both}}^{\text{pair}} - \hat{N}_{\text{one}}^{\text{pair}}$, respectively.

Comparison of experimentally verified and predicted binding sites

We define two vectors which we then compare to evaluate the distance between experimentally verified and predicted membrane binding residues. The C_α -coordinate of experimentally verified membrane binding residues functions as a proxy for the membrane, and the vector defined by the latter residues and the center of mass (COM) of the protein is used as a reference to which we compare the vector defined by the protein COM and the Likely Inserted Hydrophobe. Given a set of identified or predicted membrane interacting residues, I , we compute the vector, \mathbf{t}_I :

$$\mathbf{t}_I = \frac{1}{|I|} \sum_{a \in I} \mathbf{v}_a - \frac{1}{|G_*|} \sum_{a \in G_*} \mathbf{v}_a \tag{11}$$

where \mathbf{v}_a denotes the C_α -coordinates of residue a , and G_* is the set of all residues in the protein. We will denote vectors obtained for experimentally identified membrane binding residues as \mathbf{t}_{I_e} , and those obtained for a Likely Inserted Hydrophobe as \mathbf{t}_{I_p} . We then measure the angle $\angle \mathbf{t}_{I_e} \mathbf{t}_{I_p}$ between the two vectors for each protein in the dataset of known binding sites.

Implementation

The solvent accessible area was calculated with MMTK [69] (version 2.9.0), and the convex hull was calculated with Qhull [70] via scipy [71] (version 0.13.3). Proportion test confidence intervals were calculated with R [72] (Version 2.12.0), odds ratios and corresponding confidence intervals were calculated with the R-package epitools [73] (version 0.5-6). Secondary structure annotations were computed with the CMBI DSSP implementation [74] (version 2.0.4). Otherwise the analyses were implemented by us, using Python and R. Plots were produced with R, and other visualisations using VMD (Visual Molecular Dynamics) [75].

Supporting information

S1 Text. Supplementary tables and analysis. Analysis to assess the robustness of results to quaternary structure modelling and the definition of the reference set.

S1 Data Set. The set of peripheral membrane protein structures. 648
 Comma-separated file with PDB IDs for the set and the OPM classification of these at 649
 the time of analysis. The PDB IDs also serve to identify entries in the OPM database. 650

**S2 Data Set. The set of structures from which reference-non-binding 651
 surfaces was extracted.** Comma-separated file with PDB IDs for the set and the 652
 OPM classification of these at the time of analysis. The PDB IDs also serve to identify 653
 entries in the OPM database. 654

Acknowledgments 655

We thank Anne Gershenson at the University of Massachusetts Amherst, and Angèle 656
 Abboud at the University of Bergen for their valuable comments. This work was 657
 supported by grants from the Norwegian Research Council (FRIMEDBIO 214167 and 658
 FRIMEDBIO 251247). 659

References

1. Kutateladze TG. Translation of the phosphoinositide code by PI effectors. *Nat Chem Biol.* 2010;6(7):507–513.
2. Vögler O, Barceló JM, Ribas C, Escribá PV. Membrane interactions of G proteins and other related proteins. *Biochim Biophys Acta.* 2008;1778(7-8):1640–1652.
3. Cullen PJ. Endosomal sorting and signalling: an emerging role for sorting nexins. *Nat Rev Mol Cell Biol.* 2008;9(7):574–582.
4. Inaba T, Kishimoto T, Murate M, Tajima T, Sakai S, Abe M, et al. Phospholipase C β 1 induces membrane tubulation and is involved in caveolae formation. *Proc Natl Acad Sci USA.* 2016;113(28):7834–7839.
5. Itoh T, Erdmann KS, Roux A, Habermann B, Werner H, De Camilli P. Dynamin and the actin cytoskeleton cooperatively regulate plasma membrane invagination by BAR and F-BAR proteins. *Dev Cell.* 2005;9(6):791–804.
6. Lemmon MA. Membrane recognition by phospholipid-binding domains. *Nat Rev Mol Cell Biol.* 2008;9(2):99–111.
7. Cho W, Stahelin RV. Membrane-protein interactions in cell signaling and membrane trafficking. *Annu Rev Biophys Biomol Struct.* 2005;34(1):119–151.
8. Mulgrew-Nesbitt A, Diraviyam K, Wang J, Singh S, Murray P, Li Z, et al. The role of electrostatics in protein-membrane interactions. *Biochim Biophys Acta.* 2006;1761(8):812–826.
9. Lazaridis T. Effective energy function for proteins in lipid membranes. *Proteins: Struct, Funct, Bioinf.* 2003;52:176–192.
10. Lomize AL, Pogozheva ID, Lomize MA, Mosberg HI. The role of hydrophobic interactions in positioning of peripheral proteins in membranes. *BMC Structural Biology.* 2007;7:44.
11. Gilbert GE, Kaufman RJ, Arena AA, Miao H, Pipe SW. Four hydrophobic amino acids of the factor VIII C2 domain are constituents of both the membrane-binding and von Willebrand factor-binding motifs. *J Biol Chem.* 2002;277(8):6374–6381.

12. Gamsjaeger R, Johs A, Gries A, Gruber HJ, Romanin C, Prassl R, et al. Membrane binding of β 2-glycoprotein I can be described by a two-state reaction model: an atomic force microscopy and surface plasmon resonance study. *Biochem J*. 2005;389(3):665–673.
13. Ford MGJ, Mills IG, Peter BJ, Vallis Y, Praefcke GJK, Evans PR, et al. Curvature of clathrin-coated pits driven by epsin. *Nature*. 2002;419(6905):361–366.
14. Balali-Mood K, Bond PJ, Sansom MSP. Interaction of monotopic membrane enzymes with a lipid bilayer: A coarse-grained MD simulation study †. *Biochemistry*. 2009;48(10):2135–2145.
15. Miller S, Janin J, Lesk AM, Chothia C. Interior and surface of monomeric proteins. *J Mol Biol*. 1987;196(3):641–656.
16. Perisic O, Fong S, Lynch DE, Bycroft M, Williams RL. Crystal structure of a calcium-phospholipid binding domain from cytosolic phospholipase A2. *J Biol Chem*. 1998;273(3):1596–1604.
17. Bravo J, Karathanassis D, Pacold CM, Pacold ME, Ellson CD, Anderson KE, et al. The crystal structure of the PX domain from p40(phox) bound to phosphatidylinositol 3-phosphate. *Mol Cell*. 2001;8(4):829–839.
18. Scott DL, White SP, Otwinowski Z, Yuan W, Gelb MH, Sigler PB. Interfacial catalysis: the mechanism of phospholipase A2. *Science*. 1990;250(4987):1541–1546.
19. Zhang G, Kazanietz MG, Blumberg PM, Hurley JH. Crystal structure of the cys2 activator-binding domain of protein kinase C delta in complex with phorbol ester. *Cell*. 1995;81(6):917–924.
20. Misra S, Hurley JH. Crystal structure of a phosphatidylinositol 3-phosphate-specific membrane-targeting motif, the FYVE domain of Vps27p. *Cell*. 1999;97(5):657–666.
21. Agasøster AV, Halskau Ø, Fuglebakk E, Frøystein NA, Muga A, Holmsen H, et al. The interaction of peripheral proteins and membranes studied with alpha-lactalbumin and phospholipid bilayers of various compositions. *J Biol Chem*. 2003;278(24):21790–21797.
22. Lomize AL, Pogozheva ID, Mosberg HI. Anisotropic solvent model of the lipid bilayer. 1. Parameterization of long-range electrostatics and first solvation shell effects. *J Chem Inf Model*. 2011;51(4):918–929.
23. Lomize MA, Pogozheva ID, Joo H, Mosberg HI, Lomize AL. OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res*. 2012;40(Database issue):D370–6.
24. Wimley WC, White SH. Experimentally determined hydrophobicity scale for proteins at membrane interfaces. *Nat Struct Biol*. 1996;3(10):842–848.
25. Wilmot CM, Thornton JM. Analysis and prediction of the different types of beta-turn in proteins. *J Mol Biol*. 1988;203(1):221–232.
26. Park MJ, Sheng R, Silkov A, Jung DJ, Wang ZG, Xin Y, et al. SH2 domains serve as lipid-binding modules for pTyr-signaling proteins. *Mol Cell*. 2016;62(1):7–20.

27. Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline state. *J Mol Biol.* 2007;372(3):774–797.
28. Fox NK, Brenner SE, Chandonia JM. SCOPe: Structural classification of proteins—extended, integrating SCOP and ASTRAL data and classification of new structures. *Nucleic Acids Res.* 2014;42(Database issue):D304–9.
29. Lomize AL, Pogozheva ID, Mosberg HI. Anisotropic solvent model of the lipid bilayer. 2. Energetics of insertion of small molecules, peptides, and proteins in membranes. *J Chem Inf Model.* 2011;51(4):930–946.
30. Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, et al. The Protein Data Bank. *Nucleic Acids Res.* 2000;28(1):235–242.
31. Velankar S, Best C, Beuth B, Boutselakis CH, Cobley N, Sousa Da Silva AW, et al. PDBe: Protein Data Bank in Europe. *Nucleic Acids Res.* 2010;38(Database issue):D308–17.
32. Hedin EMK, Høystrup P, Patkar SA, Vind J, Svendsen A, Fransson L, et al. Interfacial orientation of thermomyces lanuginosa Lipase on phospholipid vesicles investigated by electron spin resonance relaxation spectroscopy. *Biochemistry.* 2002;41(48):14185–14196.
33. Grauffel C, Yang B, He T, Roberts MF, Gershenson A, Reuter N. Cation- π interactions as lipid-specific anchors for phosphatidylinositol-specific phospholipase C. *J Am Chem Soc.* 2013;135(15):5740–5750.
34. Malmberg NJ, Van Buskirk DR, Falke JJ. Membrane-docking loops of the cPLA2 C2 domain: detailed structural analysis of the protein-membrane interface via site-directed spin-labeling. *Biochemistry.* 2003;42(45):13227–13240.
35. Stahelin RV, Burian A, Bruzik KS, Murray D, Cho W. Membrane binding mechanisms of the PX domains of NADPH oxidase p40phox and p47phox. *J Biol Chem.* 2003;278(16):14469–14479.
36. Stahelin RV, Long F, Diraviyam K, Bruzik KS, Murray D, Cho W. Phosphatidylinositol 3-phosphate induces the membrane penetration of the FYVE domains of Vps27p and Hrs. *J Biol Chem.* 2002;277(29):26379–26388.
37. Wang QJ, Fang TW, Nacro K, Marquez VE, Wang S, Blumberg PM. Role of hydrophobic residues in the C1b domain of protein kinase C delta on ligand and phospholipid interactions. *J Biol Chem.* 2001;276(22):19580–19587.
38. Stahelin RV, Long F, Peter BJ, Murray D, De Camilli P, McMahon HT, et al. Contrasting membrane interaction mechanisms of AP180 N-terminal homology (ANTH) and epsin N-terminal homology (ENTH) Domains. *J Biol Chem.* 2003;278(31):28993–28999.
39. Stahelin RV, Cho W. Differential Roles of Ionic, Aliphatic, and Aromatic Residues in Membrane–Protein Interactions: A surface plasmon resonance study on phospholipases A2. *Biochemistry.* 2001;40(15):4672–4678.
40. Frazier AA, Roller CR, Havelka JJ, Hinderliter A, Cafisco DS. Membrane-bound orientation and position of the synaptotagmin I C2A domain by site-directed spin labeling. *Biochemistry.* 2003;42:96–105.

41. Gerber SH, Rizo J, Sudhof TC. Role of electrostatic and hydrophobic interactions in Ca²⁺-dependent phospholipid binding by the C2A-domain from synaptotagmin I. *Diabetes*. 2002;51(Supplement 1):S12–S18.
42. Rufener E, Frazier AA, Wieser CM, Hinderliter A, Cafiso DS. Membrane-bound orientation and position of the synaptotagmin C2B domain determined by site-directed spin labeling. *Biochemistry*. 2005;44(1):18–28.
43. Corbalán-García S, Sánchez-Carrillo S, García-García J, Gómez-Fernández JC. Characterization of the membrane binding mode of the C2 domain of PKC ϵ . *Biochemistry*. 2003;42(40):11661–11668.
44. Kim SW, Quinn-Allen MA, Camp JT, Macedo-Ribeiro S, Fuentes-Prior P, Bode W, et al. Identification of functionally important amino acid residues within the C2-domain of human Factor V using alanine-scanning mutagenesis. *Biochemistry*. 2000;39(8):1951–1958.
45. Feng J, Wehbi H, Roberts MF. Role of tryptophan residues in interfacial binding of phosphatidylinositol-specific phospholipase C. *J Biol Chem*. 2002;277(22):19867–19875.
46. Feng J, Bradley WD, Roberts MF. Optimizing the interfacial binding and activity of a bacterial phosphatidylinositol-specific phospholipase C. *J Biol Chem*. 2003;278(27):24651–24657.
47. Walther M, Wiesner R, Kuhn H. Investigations into calcium-dependent membrane association of 15-lipoxygenase-1. Mechanistic roles of surface-exposed hydrophobic amino acids and calcium. *J Biol Chem*. 2004;279(5):3717–3725.
48. Kohout SC, Corbalán-García S, Gómez-Fernández JC, Falke JJ. C2 domain of protein kinase C α : Elucidation of the membrane docking surface by site-directed fluorescence and spin labeling. *Biochemistry*. 2003;42(5):1254–1265.
49. Goh BC, Wu H, Rynkiewicz MJ, Schulten K, Seaton BA, McCormack FX. Elucidation of lipid binding sites on lung surfactant protein A using X-ray crystallography, mutagenesis, and molecular dynamics simulations. *Biochemistry*. 2016;55(26):3692–3701.
50. B Campos, Y D Mo, T R Mealy, C W Li, M A Swairjo, C Balch, et al. Mutational and crystallographic analyses of interfacial residues in annexin V suggest direct interactions with phospholipid membrane components. *Biochemistry*. 1998;37(22):8004–8010.
51. Isas JM, Langen R, Hubbell WL, Haigler HT. Structure and dynamics of a helical hairpin that mediates calcium-dependent membrane binding of annexin B12. *J Biol Chem*. 2004;279(31):32492–32498.
52. Kutateladze T, Overduin M. Structural mechanism of endosome docking by the FYVE domain. *Science*. 2001;291(5509):1793–1796.
53. Anderluh G, Razpotnik A, Podlesek Z, Macek P, Separovic F, Norton RS. Interaction of the eukaryotic pore-forming cytotoxin equinatoxin II with model membranes: 19F NMR studies. *J Mol Biol*. 2005;347(1):27–39.
54. Shenkarev ZO, Nadezhdin KD, Sobol VA, Sobol AG, Skjeldal L, Arseniev AS. Conformation and mode of membrane interaction in cyclotides. Spatial structure of kalata B1 bound to a dodecylphosphocholine micelle. *Febs J*. 2006;273(12):2658–2672.

55. Lin Y, Nielsen R, Murray D, Hubbell WL, Mailer C, Robinson BH, et al. Docking phospholipase A2 on membranes using electrostatic potential-modulated spin relaxation magnetic resonance. *Science*. 1998;279(5358):1925–1929.
56. Canaan S, Nielsen R, Ghomashchi F, Robinson BH, Gelb MH. Unusual mode of binding of human group IIA secreted phospholipase A2 to anionic interfaces as studied by continuous wave and time domain electron paramagnetic resonance spectroscopy. *J Biol Chem*. 2002;277(34):30984–30990.
57. Lathrop B, Gadd M, Biltonen RL, Rule GS. Changes in Ca²⁺ affinity upon activation of *Agkistrodon piscivorus piscivorus* phospholipase A2. *Biochemistry*. 2001;40(11):3264–3272.
58. Chen X, Wolfgang DE, Sampson NS. Use of the parallax-quench method to determine the position of the active-site loop of cholesterol oxidase in lipid bilayers. *Biochemistry*. 2000;39(44):13383–13389.
59. Sekino-Suzuki N, Nakamura M, Mitsui KI, Ohno-Iwashita Y. Contribution of individual tryptophan residues to the structure and activity of theta-toxin (perfringolysin O), a cholesterol-binding cytolysin. *Eur J Biochem*. 1996;241(3):941–947.
60. Phillips LR, Milescu M, Li-Smerin Y, Mindell JA, Kim JI, Swartz KJ. Voltage-sensor activation with a tarantula toxin as cargo. *Nature*. 2005;436(7052):857–860.
61. Thennarasu S, Lee DK, Poon A, Kawulka KE, Vederas JC, Ramamoorthy A. Membrane permeabilization, orientation, and antimicrobial mechanism of subtilisin A. *Chem Phys Lipids*. 2005;137(1-2):38–51.
62. Tatulian SA, Qin S, Pande AH, He X. Positioning membrane proteins by novel protein engineering and biophysical approaches. *J Mol Biol*. 2005;351(5):939–947.
63. Oldham ML, Brash AR, Newcomer ME. Insights from the X-ray crystal structure of coral 8R-lipoxygenase: calcium activation via a C2-like domain and a structural basis of product chirality. *J Biol Chem*. 2005;280(47):39545–39552.
64. Mathias JD, Ran Y, Carter JD, Fanucci GE. Interactions of the GM2 activator protein with phosphatidylcholine bilayers: A site-directed spin-labeling power saturation study. *Biophys J*. 2009;97(5):1436–1444.
65. Jian X, Tang WK, Zhai P, Roy NS, Luo R, Gruschus JM, et al. Molecular basis for cooperative binding of anionic phospholipids to the PH domain of the Arf GAP ASAP1. *Structure*. 2015;23(11):1977–1988.
66. Eisenhaber F, Lijnzaad P, Argos P, Sander C, Scharf M. The double cubic lattice method: Efficient approaches to numerical integration of surface area and volume and to dot surface contouring of molecular assemblies. *J Comput Chem*. 1995;16(3):273–284.
67. Bondi A. van der Waals volumes and radii. *J Phys Chem*. 1964;68(3):441–451.
68. Kabsch W, Sander C. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers*. 1983;22(12):2577–2637.
69. Hinsen K. The molecular modeling toolkit: A new approach to molecular simulations. *J Comput Chem*. 2000;21(2):79–85.

70. Barber CB, Dobkin DP, Huhdanpaa H. The quickhull algorithm for convex hulls. *ACM T Math Software*. 1996;22(4):469–483.
71. Jones E, Oliphant E, Peterson P. *SciPy: Open Source Scientific Tools for Python*. 2001;.
72. Team RDC. *R: A language and environment for statistical computing*. 2008;.
73. Aragon T. epitools: Epidemiology Tools. R package. 2010;.
74. Touw WG, Baakman C, Black J, te Beek TAH, Krieger E, Joosten RP, et al. A series of PDB-related databanks for everyday needs. *Nucleic Acids Res*. 2015;43(D1):D364–D368.
75. Humphrey W, Dalke A, Schulten K. VMD: Visual molecular dynamics. *J Mol Graph*. 1996;14(1):33–38.